

52013/CAB/A400



SUBSTITUTE SPECIFICATION  
(CLEAN COPY)

PETUNIA PLANT NAMED 'SUNPURPLE'

BOTANICAL CLASSIFICATION

*Petunia hybrida*

VARIETAL DENOMINATION

5 'Sunpurple'

BACKGROUND OF THE VARIETY

The present invention relates to a new and distinct variety of Petunia plant, which originated from crossing a seedling of a Petunia variety called 'Red Madness'  
10 (unpatented) as the female parent and a Petunia wild species called '70-200' (unpatented) as the male parent.

The Petunia is a very popular plant that is used for flower bedding and potting in the summer season. There are only a few Petunia varieties that do not have an upright  
15 growth habit and that have a high resistance to rain, heat, and diseases. The Petunia plants such as REVOLUTION series, 'Revolution Purple pink' (U.S. Plant Patent No. 6,915), 'Revolution Brilliant pink' (U.S. Plant Patent No. 6,914), 'Revolution Brilliantpink-Mini' (U.S. Plant Patent No. 6,899),  
20 and 'Revolution Blue vein' (U.S. Plant Patent No. 9,322) are decumbent type plants having long stems, a lower plant height, abundant branching, and a high resistance to heat, rain and diseases. However, there are only a few Petunia varieties having a decumbent plant shape, a great profusion of flowers,  
25 vivid reddish purple petals and a high resistance to rain, heat, and diseases. Accordingly, this invention was aimed at obtaining a new variety having vivid reddish purple petals together with the above features.

The female parent 'Red Madness' used in the crossing to  
30 produce 'Sunpurple' is a cultivar having compact and spreading growth habit with medium size single flowers, the petals

52013/CAB/A400

having a strong red color.

The male parent '70-200' used in the crossing to produce 'Sunpurple' is a strain of wild Petunia species having decumbent growth habit with much branching. It has small size  
5 single flowers, the petals having vivid reddish purple color.

In April 2000, crossing of 'Red Madness' as the female parent and '70-200' as the pollen parent was conducted at Yokaichi, Shiga, Japan. In August 2000, 80 seedlings were obtained from that crossing. These seedlings were grown in  
10 pots in glasshouses and were evaluated. One seedling was selected in view of its growth habit, flower size and color in October 2000. That seedling was propagated by cuttings and a trial was carried out by flower potting and bedding from April to September 2001. The botanical characteristics of that  
15 plant were then examined, using similar varieties 'Sunripami' (unpatented) and 'Sunrovein' (U.S. Plant Patent Application Serial No. 10/789,408) for comparison. As a result, it was concluded that this Petunia plant is distinguishable from any other variety, whose existence is known to us, and uniform and  
20 stable in its characteristics. The instant plant reproduces true to type in successive generations of asexual reproduction. The new variety of Petunia plant was named 'Sunpurple'.

In the following description, the color-coding is in  
25 accordance with the Colour Chart of The Royal Horticultural Society, London, England (R.H.S.).

#### SUMMARY OF THE VARIETY

This new variety is unlike any Petunia commercially available variety known to the inventors, as evidenced by the  
30 following unique combinations of characteristics.

1. Decumbent growth habit with long stems.
2. Abundant branching and great profusion of blooms.
3. The flowers are single and medium size. The petal

52013/CAB/A400

color is vivid reddish purple (near R.H.S. N74A).

4. The plant has a high resistance to cold, heat, rain and diseases.

The new variety 'Sunpurple' differs from the similar variety 'Sunripami' in the following points.

1. The leaf of 'Sunpurple' is longer than that of 'Sunripami'.

2. The leaf of 'Sunpurple' is thinner than that of 'Sunripami'.

3. The flower of 'Sunpurple' is larger than that of 'Sunripami'.

4. The bottom color of the corolla throat of 'Sunpurple' is deep reddish purple (near R.H.S. 72A), while that of 'Sunripami' is moderate purple (near R.H.S. 83B).

5. The outside color of the corolla throat of 'Sunpurple' is deep reddish purple (near R.H.S. N79B), while that of 'Sunripami' is strong purple (near R.H.S. 83D).

6. The apex shape of the petal tip of 'Sunpurple' is obtuse. That of 'Sunripami' is rounded.

7. The flowering time of 'Sunpurple' is later than that of 'Sunripami'.

The new variety 'Sunpurple' differs from the similar variety 'Sunrovein' in the following points.

1. The plant height of 'Sunpurple' is lower than that of 'Sunrovein'.

2. The leaf of 'Sunpurple' is longer than that of 'Sunrovein'.

3. The leaf of 'Sunpurple' is thinner than that of 'Sunrovein'.

4. The petal color of 'Sunpurple' is vivid reddish purple (near R.H.S. N74A), while that of 'Sunrovein' is vivid purplish red (near R.H.S. 71B) with deep purplish red (near

52013/CAB/A400

R.H.S. 71A) vein.

5        5.     The bottom color of the corolla throat of  
'Sunpurple' is deep reddish purple (near R.H.S. 72A), while  
that of 'Sunrovein' is moderate purplish red (near R.H.S.  
64A).

6.     The outside color of the corolla throat is deep  
reddish purple (near R.H.S. N79B), while that of 'Sunrovein'  
is light purplish pink (near R.H.S. 62C).

#### BRIEF DESCRIPTION OF THE PHOTOGRAPH

10        The depicted plants had been produced by the use of  
cuttings and were photographed during July 2003 at an age of  
approximately 6 months while growing outdoors in 15 cm pots at  
Yokaichi-shi, Shiga-ken, Japan, under typical outdoor  
conditions for that region.

15        FIG. 1 is a photograph of a typical plant of the new  
variety of Petunia plant 'Sunpurple'.

FIG. 2 is a photograph of flowers and leaves of the new  
variety of Petunia plant 'Sunpurple'.

#### DESCRIPTION OF THE VARIETY

20        The botanical characteristics of the new and distinct  
variety of Petunia plant named 'Sunpurple' at an age of 4  
months, at Yokaichi, Shiga, Japan, are as follows.

Plant:

Growth habit. - Decumbent.

25        Plant height. - Approximately 12.3 cm.

Spreading area of plant. - Approximately 33.2 cm.

Blooming period. - April to late October in the southern  
Kanto area, Japan. The plant shape does not change throughout  
this period. A typical flower commonly lasts approximately 5  
30        days on the plant when experiencing a temperature of  
approximately 20°C.

Stem:

52013/CAB/A400

Length. - Approximately 30 cm.

Thickness. - Approximately 2.5 mm.

Pubescence. - Present and typical for the species.

Branching. - Abundant.

5 Internode length. - Approximately 2.2 cm.

Color. - Near R.H.S. 144A (strong yellow green).

Leaf:

Whole shape. - Elliptic. The apex shape is acute, and the base shape is attenuate.

10 Margin. - Entire.

Length. - Approximately 6.4 cm.

Width. - Approximately 3.5 cm.

Color. - Upper side color is near R.H.S. 144A (strong yellow green). Bottom side color is near R.H.S. 146C (moderate olive green).

15

Thickness. - Approximately 0.2 mm.

Pubescence. - Sparse.

Petiole: (Indistinct)

Length. - Approximately 3.0 mm.

20 Diameter - Approximately 1.0 mm.

Color. - Near R.H.S. 144B.

Buds:

Shape. - Cylindrical.

Length. - Approximately 4.5 cm.

25 Diameter. - Approximately 7.0 mm.

Color. - Toward apex, Near R.H.S. N77B; base, near R.H.S. N79B; venation, near R.H.S. N92A.

Flower:

Facing direction. - Slanted upward.

30 Type. - Single.

Shape. - Funnel-shape, with five-fissured limb.

Diameter. - Approximately 7.4 cm.

Depth. - Approximately 5.8 cm.

52013/CAB/A400

Throat diameter. - Distal end, approximately 1.5 cm.

Tube Diameter. - Proximal end, approximately 4.0 mm.

Petals:

Whole shape. - Very broadly obtrullate.

5 Apex shape. - Obtuse.

Width. - Approximately 4.0 cm.

Length. (from the throat). - Approximately 3.5 cm.

Margin. - Entire, undulated.

10 Color. - Upper side of the petal, near R.H.S. N74A;  
underside of the petal, near R.H.S. N78A with near N92A  
venation.

Upper color of the corolla throat. - Near R.H.S. N90B  
with N92A venation.

Bottom color of the corolla throat. - Near R.H.S. 72A.

15 Outside color of corolla tube. - Near R.H.S. N79B.

Reproductive organs. - 1 normal pistil and 5 normal  
stamens. Color of pistil is near R.H.S. 150C (light yellow  
green). Color of stamen is near R.H.S. 157C (light yellow  
green).

20 Peduncle. - Approximately 1.3 mm in diameter and approximately  
1.9 cm in length.

Color. - Near R.H.S. 144B.

Surface. - Pubescent.

Calyx. - Medium. 5 sepals fused at the base.

25 Sepals:

Shape. - Narrow elliptic.

Apex shape. - Rounded.

Base. - Fused.

Margin. - Entire, undulated.

30 Surface. - Pubescent.

Length. - Approximately 2.5 cm.

Width. - Approximately 6.0 mm.

Color. - Upper surface, near R.H.S. 144A; lower surface,

52013/CAB/A400

toward apex, near R.H.S. 144A; base, near R.H.S. N77A.

Physiological and ecological characteristics.- High resistance to cold, heat, rain and diseases. Moderate resistance to pests.

- 5        This new variety of Petunia plant is most suitable for flower bedding and potting, particularly in hanging pots or planters, and is excellent for use as ground cover. Pinching of old blossoms will enhance the formation of new blossoms.